

TY SABRI INFUSION ORDERS

Patient Name: _____

Diagnosis: _____

***** Due to the risk of Progressive Multifocal Leukoencephalopathy (PML), Tysabri is restricted through a "REMS" requirement and is only available through the "TOUCH" Prescribing Program. THE NURSE MUST CONFIRM PATIENT IS AUTHORIZED to receive Tysabri; Via on-line portal. Complete Pre-Infusion Patient Checklist & provide patient with the Medication Guide. As part of the REMS requirement, the provider must authorize continued treatment every 6 months.**

- * Screen patients at each visit for active infection before any treatment is given, especially for any new signs/symptoms suggestive of PML (e.g. changes in thinking, memory, eyesight, balance, or strength), encephalitis, or meningitis (e.g., fever, H/A, confusion). If present, notify provider immediately.
- * Screen patients for pregnancy status and notify provider for possible pregnancy.
- * Notify provider of any signs/ symptoms suggestive of liver disease (e.g., jaundice, dark urine, nausea, abdominal pain, fatigue).
- * Avoid use of 'live' vaccinations concurrently with Tysabri.

The following orders will be enacted unless a specific order is written to the contrary:

1. Please check to activate each order desired:

- Natalizumab (Tysabri) 300 mg in 100 ml 0.9% Sodium Chloride IV infused over 1 hour every 4 weeks. Following infusion flush line with at least 20 ml of 0.9% Sodium Chloride IV.

2. Monitor patient for hypersensitivity reaction during and for at least 1 hour after infusion completed.

3. Additional meds/IV fluids (Tysabri compatible with N/S only): _____

4. Premedication: If patient experiences an infusion-related reaction, consider pre-medication prior to subsequent infusions. Patients who have experienced *hypersensitivity* reactions should **NOT** be retreated with Tysabri.

<input type="checkbox"/> Acetaminophen (Tylenol) 500mg PO X 1	<input type="checkbox"/> Patient instructed to take at home
Loratadine (Claritin) 10mg PO X 1 --OR--	<input type="checkbox"/> Patient instructed to take at home
<input type="checkbox"/> Diphenhydramine (Benadryl) 50mg PO X 1	
<input type="checkbox"/> Prednisone _____ mg PO PM night before infusion	<input type="checkbox"/> Patient instructed to take at home
Prednisone _____ mg PO AM morning of infusion	<input type="checkbox"/> Patient instructed to take at home

5. Infusion Reaction protocol: The following orders will be enacted unless a specific order is written to the contrary:

For **MINOR** infusion reaction (e.g., headache, fever, chills, nausea, flushing)

- Continue infusion at current or decreased rate as tolerated by patient; monitor patient closely

For **MODERATE** infusion reaction (e.g., pruritus, rash, dizziness)

- STOP infusion
- Give diphenhydramine (Benadryl) 25mg IV x 1 dose
- May restart infusion after 15 minutes at a decreased rate as tolerated only if patient is asymptomatic and vital signs are stable.
Notify provider

For **SEVERE** infusion reaction or anaphylaxis (e.g., urticaria, dyspnea, wheezing, hypotension, abdominal cramps, angioedema)

- **STOP administration of Tysabri immediately**
- For ANAPHYLAXIS: Epinephrine (EpiPen) 0.3mg (0.3ml) IM x 1 STAT, administered into anterolateral aspect of the thigh
- For HYPOTENSION: Bolus IV 0.9% Sodium Chloride 1000 ml over 1 hour
- Diphenhydramine (Benadryl) 25 mg IV X 1 dose
Methylprednisolone (Solu-Medrol) 125 mg IV x 1 dose
- Notify provider
- Transport the patient to the Emergency Department

PHYSICIAN SIGNATURE

DATE/TIME