



Patient Name: _____

DOB: _____

UPLIZNA® (INEBILIZUMAB-CDON) ORDER SET

Diagnosis:

G36.0 Neuromyelitis Optica Spectrum Disorder _____ Other: _____
(ICD-10)

Prescriber must indicate all of the following requirements have been met (attach supporting documentation):

- quantitative immunoglobulins within normal limits
 anti-aquaporin-4 (AQP4) antibody positive (*required*)
 Latent TB screening **negative**
 HBV screening **negative**

If any the above are *not* checked, attach treatment/consultation notes clearing patient for inebilizumab-cdon therapy

Pre-Infusion:

- Assess for contraindications; hold infusion and notify provider for:**
- signs/symptoms of active infection;
 - planned or recent invasive/surgical procedure;
 - receipt of live or live-attenuated vaccines within 4 weeks;
 - chance of pregnancy; or
 - signs/symptoms of PML (new or worsening unilateral weakness, confusion or changes in vision, thinking, memory, balance, or personality/mood).
- Obtain vital signs at baseline and with rate changes Establish vascular access
- If infusion-related reaction occurs, stop infusion and treat per orders/protocol as clinically indicated.

Pre-medications: (Prescriber must select *one* option within each set of brackets for each medication):

- acetaminophen [500 mg 650 mg] PO once [30 60] min prior to infusion
 methylprednisolone [80 mg 125 mg _____ mg] IV once [30 60] min prior to infusion
 diphenhydramine [25 mg 50 mg _____ mg] [IV PO] once [30 60] min prior to infusion

Medication Orders:

- Dilute **inebilizumab-cdon 300 mg/30 mL in 250 mL 0.9% sodium chloride** and administer intravenously using a sterile, in-line, low protein-binding **0.2- or 0.22-micron filter** using rates in table at right.

Elapsed Time (minutes)	Infusion Rate
0-30	42 mL/hr
31-60	125 mL/hr
61 to completion	333 mL/hr

Post-Infusion:

- Flush administration set with 0.9% sodium chloride to deliver residual volume.
- Record vital signs immediately following infusion and prior to discharge.
- Leave IV in place for observation period; remove prior to discharge.
- Monitor patient for hypersensitivity reaction for a period of 60 minutes following infusion.**
- Provide patient with discharge instructions.
- Send record of treatment to prescriber at fax number below.
- Follow standing anaphylaxis order and notify MD of outcome.

Frequency:

- On Day 1 and Day 15; repeat in 6 months (from Day 1) Every 6 months (date of last treatment: _____)

Additional Orders:

Prescriber Name (print): _____ Fax: _____

Prescriber signature: _____

Date: _____

Fax Order To: 888-360-2455

Order valid for one year unless otherwise indicated.