

RHEUMATOLOGY ORDER

PATIENT INFORMATION

TREATMENT ARRANGEMENTS

Last Name _____ First Name _____
 Social Security # _____ Date of Birth _____
 Address: _____
 Sex M F Phone: _____

Start Date: _____
 Ship Meds: Home Doctor Office
 Other: _____

Diagnosis: Rheumatoid Arthritis Ankylosing Spondylitis Osteoporosis Psoriatic Arthritis Lupus Ulcerative Colitis Gout Crohn's
 Spondyloarthropathy and/or Other: _____ Date of Diagnosis: _____

Premeds: Tylenol 500mg 2 PO Loratadine 10mg PO Benadryl _____ mg IV/PO (circle one) SoluCortif _____ mg IV
 Ondansetron _____ mg IV Promethazine _____ mg IV Quzyttir 10mg IVSP over 1-2 min Other SoluMedol _____ mg IV

√ Standing Order: Anaphylaxis Protocol √ **Skilled Nurse to start PIV, infuse per protocol, DC PIV each visit**

Lab Draw As Follows: _____ Q _____
 Quantiferon Gold Lab Draw Q _____ Patient to FU w/MD Q _____

PRESCRIPTION INFORMATION

MEDICATION	STRENGTH	DIRECTIONS	QTY	REFIL
<input type="checkbox"/> Actemra	<input type="checkbox"/> 80 mg/4 ml <input type="checkbox"/> 200 mg/10 ml _____ mg/kg <input type="checkbox"/> 400 mg/20 ml	Induction dose: 4 mg/kg every 4 weeks Maint. Dose: (based on clinical response): 8 mg/kg every 4 weeks Other:		
<input type="checkbox"/> Avsola	5mg/kg	Induction Dose: Administer IV on week 0, week 2, week 6 Maint. Dose: IV every 8 weeks after Induction dose		
<input type="checkbox"/> Benlysta	40 mg Vial / 10mg/kg	Induction Dose: Administer IV over 1 hour on week 0, then q2w x 3 doses Maint. Dose: IV over 1 hour every 4 weeks after Induction dose		
<input type="checkbox"/> Inflectra	3mg/kg 5mg/kg 10mg/kg	Induction Dose: Administer IV over 1 hour on week 0, week 2, week 6 Maint. Dose: IV over 1 hour every 8 weeks after Induction dose		
<input type="checkbox"/> Kyrstexxa	8mg/ml <i>*protect from light and use within 4 hours of mixing</i> Confirm uric acid level prior to infusion Methotrexate (As Prescribed - Immunomodulator)	8mg/mL IV in bag of 0.9% or 0.45%NaCl every 2 weeks over 2 hours IV Corticosteroids: Methylprednisolone hydrocortisone 125mg prior to each infusion. Antihistamines: Allegra 180mg; Claritin 10mg; Benadryl 25-50mg po to be taken night before infusion and/or can administered concomitant with infusion. Oral Analgesic: Tylenol 325mg 2 po prior to each infusion.		
<input type="checkbox"/> Orencia	250mg Vial	_____ mg in 100ml of 0.9% NaCl over 30 min. at weeks 0, 2, and 4, then every 4 weeks		
<input type="checkbox"/> Rituxan	<input type="checkbox"/> 100mg/10ml vial <input type="checkbox"/> 500mg/50ml vial	Infuse two doses of 100mg in 1 liter of 0.9% NaCl separated by 2 weeks. Other:		
<input type="checkbox"/> Remicade	_____ mg/kg 100mg Vial	Induction Dose: IV in 250ml of 0.9% NaCl at weeks 0, 2, and 6 weeks. Maint. Dose: IV in 250ml of 0.9% NaCl every 8 weeks. Maint. Dose: IV in 250ml of 0.9% NaCl every 6 weeks. Other:		
<input type="checkbox"/> Renflexis	3mg/kg 5mg/kg 100mg Vial	Induction Dose: IV in 250ml of 0.9% NaCl at weeks 0, 2, and 6 weeks. Maint. Dose: IV in 250ml of 0.9% NaCl every 8 weeks. Maint. Dose: IV in 250ml of 0.9% NaCl every 6 weeks. Other:		
<input type="checkbox"/> Ruxience	500mg Methylprednisolone 100mg IV or equivalent 30 min prior to infusion	2 - 1000mg IV infusions over 90 min separated by 2 weeks every 24 weeks Other: Every _____ weeks Induction: 375mg/per meter square once weekly for 4 weeks for active GPA or MPA. Followed by Two 500mg infusions separated by 2 weeks followed by 500mg every 6 months		
<input type="checkbox"/> Simponi Aria	<input type="checkbox"/> _____ mg/kg	Induction Dose: Administer IV over 30 min on week 0, week 4, then 8 weeks Maint. Dose: IV over 30 min every 8 weeks after Induction dose		
<input type="checkbox"/> Saphnelo	300mg	IV 30 min every 4 weeks for _____ months		
<input type="checkbox"/> Truxima	100mg/10mL / 500mg/50mL 1000mg	Induction Dose: 50mg/hr, increase rate by 50mg/hr every 30 min, max rate 400mg/hr Maint. Dose: 100mg/hr, increase rate by 100mg/hr every 30 min, max rate 400mg/hr		
<input type="checkbox"/> OMVOH or other	300mg/15mL other:	INITIAL: administered by IV over at least 30 minutes at weeks 0, 4, and 8 MAINTENANCE: 200mg administered by SQ, given as 2 consecutive injections of 100mg each at week 12 & every 4 weeks thereafter other:		

Skilled Nursing visit for self-injection training and one additional visit with next dose if needed

Physician Signature: _____ DAW (Dispense as Written)

Date: _____

Physician Address: _____

Phone: _____ Fax: _____

Phone: _____ Phone: _____

Physicians Name

<input type="checkbox"/>	<input type="checkbox"/>
<input type="checkbox"/>	<input type="checkbox"/>
<input type="checkbox"/>	<input type="checkbox"/>
<input type="checkbox"/>	<input type="checkbox"/>

NPI: _____

DEA No. _____

IMPORTANT NOTICE: This message may contain privileged and confidential information and is intended only for the individual named. If you are not the name addressed, you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document by mistake. Then destroy this document. Please direct all verification or notification to the company listed below.

