CURBSIDE INFUSION SERVICES

Gastroenterology Referral Form

of Pages Faxed: _____
Fax Referral To: 281-406-1047
Phone: 281-406-1046

Date Required: _____ Ship To: ☐ Patient ☐ MD Office ☐ Other: _____

PATIENT INFORMATION

Patient Name: _____
Address: _____
City, State, Zip: _____
Home Phone: _____
Cell Phone: _____
Date of Birth: _____ ☐ Male ☐ Female
Emergency Contact: _____ Phone: _____

PRESCRIBER INFORMATION

Prescriber Name: _____
Address: _____
City, State, Zip: _____
Phone: _____
Fax: _____
DEA #: _____ NPI #: _____
Contact Person: _____

INSURANCE INFORMATION (Please attach the front and back of insurance and prescription drug card.)

Primary Insurance: _____ ID: _____ Group: _____
Secondary Insurance: _____ ID: _____ Group: _____
Prescription Card: _____ ID: _____ BIN: _____ PCN: _____

To better serve your patient and facilitate insurance authorization, please complete the pertinent sections:

PATIENT DIAGNOSIS/CLINICAL INFORMATION

☐ K50.00 Crohn's Disease
☐ K51.90 Ulcerative Colitis
☐ Other: _____

Prior Medication Failed: _____
Length of Treatment: _____
Reason for Discontinuation: _____

TB/PPD test: ☐ Positive ☐ Negative ☐ Date Read: _____
Weight: _____ ☐ kg ☐ lbs Height: _____ ☐ cm ☐ in %BSA: _____
Allergies: _____ ☐ NKDA
Injection Training/Home Health RN visit is necessary. ☐ Yes ☐ No
Site of Care: ☐ Home ☐ MD Office ☐ Other: _____

PRESCRIPTION INFORMATION

Medication:	Dose/Strength:	Directions:	Refills:
☐ Cimzia	☐ 200 mg prefilled syringe ☐ 200 mg vial	☐ **INITIAL:** Inject 400 mg (two 200 mg injections) SQ on day 0, 14, and 28 (Quantity: 6) ☐ **MAINTENANCE:** Inject 400 mg (two 200 mg injections) SQ every 4 weeks (Quantity: 2)	
☐ Entyvio	☐ 300 mg vial	☐ **INITIAL:** Infuse 300 mg IV over 30 minutes at day 0, 14, and 42 (Quantity: 3) ☐ **MAINTENANCE:** Infuse 300 mg IV over 30 minutes every _____ weeks (Quantity: 1)	
☐ Monoferric	☐ Weight 50kg or more ☐ Weight Less Than 50kg	☐ 1000mg IV x 1 Dose ☐ 20mg/kg IV x 1 Dose	
☐ Inflectra ☐ Remicade ☐ Renflexis	☐ 100 mg vial	☐ **INITIAL:** Infuse IV _____ mg/kg (Dose _____mg) at 0, 2, and 6 weeks (Quantity: _____) ☐ **MAINTENANCE:** Infuse IV _____ mg/kg (Dose _____ mg) every _____ weeks (Quantity: _____) ☐ Other: _____ ☐ Round to the nearest 100 ☐ Give exact dose (do NOT round)	
☐ Omvoh	☐ Ulcerative Colitis	☐ **INITIAL:** Week 0, Week 4, and Week 8: Infuse 300mg IV over 30 minutes ☐ **MAINTENANCE:** Week12 and every 4 weeks thereafter: Inject 200mg sq given as 1 injection of 200mg or 2 consecutive injections of100mg sq	
☐ Omvoh	☐ Crohn's	☐ **INITIAL**: Week 0, Week 4 and Week 8: Infuse 900mg IV over 90 minutes. **MAINTENANCE**: Week 12 and every 4 weeks thereafter Inject 300mg sq	
☐ Stelara	☐ 130 mg/26mL vials ☐ 90 mg (2x 45 mg vials)	☐ **INITIAL:** Weight based dosing, infuse IV up to 55 kg = 260 mg (2 vials), > 55 kg to 85 kg = 390 mg (3 vials), > 85 kg = 520 mg (4 vials) ☐ **MAINTENANCE:** Inject 90 mg SQ 8 weeks after initial dose, then every 8 weeks thereafter	
☐ Other:			

Pre-Medications & Other Medications
► Infusion supplies as per protocol
► Anaphylaxis Kit as per protocol

☐ Acetaminophen _____ mg PO prior to infusion
☐ Diphenhydramine _____ mg ☐ PO ☐ IV
☐ 250ml 0.9% NaCl for hydration
☐ Quzyttir 10mg IVSP over 1-2 minutes
Other:

Flush Protocol
► NaCl 0.9% 10ml
► Before and after infusion

By signing this form and using our services, you are authorizing Curbside Infusion to serve as your prior authorization designated agent in dealing with prescription and medical insurance companies.

Prescriber Signature: _____ **Date:** _____